



Your Title Goes Here: A Subtitle May Be Included*

1stAuthor Name^{1,†}, 2ndAuthor Name^{2,‡}, and 3rdAuthor Name³

¹Department of Mathematics, Faculty of Science
Chiang Mai University, Chiang Mai 50200, Thailand

²School of Mathematics, Institute of Science
Suranaree University of Technology, Nakhon Ratchasima 30000, Thailand

³Department of Mathematics and Computer Science, Faculty of Science
Chulalongkorn University, Bangkok 10330, Thailand

Abstract

Please write an abstract of no more than 200 words here. The abstract should be sufficiently informative so as to raise the participants' interest in your work. A motivating abstract will have the following structure: 1) Background: Place the question addressed in a broad context and highlight the purpose of the study; 2) Methods: Describe briefly the main methods or treatments applied; 3) Results: Summarize the article's main findings; and 4) Conclusions: Indicate the main conclusions or interpretations. The abstract should be an objective representation of the article, it must not contain results which are not presented and substantiated in the main text and should not exaggerate the main conclusions.

Keywords: keyword 1, keyword 2, keyword 3, keyword 4, keyword 5.

2020 MSC: Primary xxXxx; Secondary xxXxx, xxXxx, xxXxx.

1 Introduction

Your text goes here. Separate text sections with the standard L^AT_EX sectioning commands. Let's do some mathematics. Let us briefly remark on the notation used throughout. Symbols \mathbf{x}, \mathbf{y} will denote vectors in Euclidean space written as column vectors, while the Greek symbol γ will denote a row vector. Consequently, \mathbb{R}^n will denote the space of column vectors and $\widehat{\mathbb{R}^n}$ that of row vectors. Whenever this distinction imposes inconveniences, for example in arguments of functions, elements of \mathbb{R}^n may be expressed as row vectors as well. The inner product in \mathbb{R}^n is $\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{y}^T \mathbf{x}$, and similarly, the duality between \mathbb{R}^n and $\widehat{\mathbb{R}^n}$ will be denoted by $\langle \mathbf{x}, \gamma \rangle = \gamma \mathbf{x}$. Integrals with respect to the Haar measure of a group H will be denoted by dh , and Δ_H will denote the modular function on H . For further details, see [1].

*This research was financially supported by ... [optional text].

[†]Speaker. [‡]Corresponding author.

Email: first.author@cmu.ac.th (A. Surname), second.author@sut.ac.th (B. Surname), third.author@gmail.com (C. Surname).

2 Preliminaries

Use the \LaTeX automatism for your citations [1–4].

2.1 Subsection Heading

One may use inline equations, $y' + 4y^2 = 0$, or displayed equations

$$\vec{a} \times \vec{b} = \vec{c} + \sum_{i=1}^n C_i$$

Equations will be labeled by section with equation numbers located on the right: Consider

$$h = T \left(\sum_{i=1}^n x_i \otimes y_i \right). \quad (2.1)$$

Please note that all internal labels for equations and all cites should be prefixed by the author's name in the form: *YourName:ref*. For example, if your last name is "Peters" then

```
\begin{equation}\label{peters:eq15}
h =T \left ( \sum_{i=1}^n x_i \otimes y_i \mathbb{R} \right ).
\end{equation}
```

2.1.1 Subsubsection Heading

Your text goes here. Use the \LaTeX automatism for cross-references as well as for your citations, see Section 2.1 and also equation (2.1).

Definition 2.1. Let $A \subseteq \mathbb{R}^n$ be a convex set. A point $x \in A$ is called an extreme point if ...

Theorem 2.2. *Theorem text goes here.*

Proof. Nobody has ever been able to give a correct proof. □

Lemma 2.3. *Lemma text goes here.*

3 Main Results

Problem 3.1. Details of the problem are described here in detail¹.

Acknowledgment. (optional) The authors are grateful to the referees for their careful reading of the manuscript and their useful comments.

References

- [1] G. B. Folland, *Real analysis: modern techniques and their applications*, 2nd ed., Wiley, New York, 1999.
- [2] R. V. Moody and J. Patera, *Fast recursion formula for weight multiplicities*, Bull. Amer. Math. Soc. **7**(2) (1982), 237–242.
- [3] B. Rom and D. Walnut, *Sampling on unions of shifted lattices*, ch. Harmonic Analysis and Applications, pp. 289–323, Birkhäuser, Boston, 2006.
- [4] A. Q. Stein, *Tight frames and coding theory*, Proceedings of the 2013 Annual Meeting in Harmonic Analysis, 2013, Glasgow, August 1–4, 2013, pp. 23–42.

¹This is a regular footnote